

Complex modular architecture around a simple toolkit of wing pattern genes

Steven M. Van Belleghem^{1,2,*†}, Pasi Rastas^{3†}, Alexie Papanicolaou⁴, Simon H. Martin³, Carlos F. Arias^{2,5}, Megan A. Supple², Joseph J. Hanly³, James Mallet⁶, James J. Lewis⁷, Heather M. Hines⁸, Mayte Ruiz¹, Camilo Salazar⁵, Mauricio Linares⁵, Gilson R. P. Moreira⁹, Chris D. Jiggins³, Brian A. Counterman^{10‡}, W. Owen McMillan^{2‡} and Riccardo Papa^{1‡}

Identifying the genomic changes that control morphological variation and understanding how they generate diversity is a major goal of evolutionary biology. In *Heliconius* butterflies, a small number of genes control the development of diverse wing colour patterns. Here, we used full-genome sequencing of individuals across the *Heliconius erato* radiation and closely related species to characterize genomic variation associated with wing pattern diversity. We show that variation around colour pattern genes is highly modular, with narrow genomic intervals associated with specific differences in colour and pattern. This modular architecture explains the diversity of colour patterns and provides a flexible mechanism for rapid morphological diversification.

Recent adaptive radiations, such as the *Heliconius* butterflies¹, Galápagos finches² and African cichlids³, offer insight into evolutionary and ecological forces that underlie diversification. Typically, ecological opportunities allow natural and sexual selection to drive adaptive change and speciation. At a genetic level, recruitment from ancient polymorphism, introgression of adaptive variants between populations and *de novo* mutation are important sources of variation. However, the genetic architecture of the traits under natural and sexual selection that facilitates rapid diversification remains largely unexplored.

In this study, we sequenced the genome of the Neotropical butterfly *Heliconius erato* and used re-sequence data from 116 additional individuals to dissect the architecture of genomic variation associated with their vividly coloured wing patterns. With over 400 different wing colour forms among 46 described species⁴, *Heliconius* represents one of the most visually diverse radiations in the animal kingdom and an excellent system for establishing a broad and integrative view of morphological diversification. The evolution of scale cells and the spatial coordinate system that controls wing pigmentation is a key innovation of the Lepidoptera. Wing patterns are often under strong natural and sexual selection, and these forces probably shape much of the pattern diversity we see among the more than 160,000 butterfly and moth species⁵.

In *Heliconius*, conspicuous wing patterns are important for signalling toxicity to potential predators⁶ and play a role in mate selection⁷. Natural selection favors Müllerian mimicry among toxic butterflies, resulting in convergence between co-occurring species, as well as geographic divergence between populations of the same species⁸. Among *Heliconius* butterflies, the genetic basis of this

wing diversity has been studied for nearly 60 years and more than 30 Mendelian loci have been described⁹. Over the past decade, however, genetic research has shown that most of the complexity of colour variation across *Heliconius* is actually controlled by relatively few genes acting broadly across the fore- and hindwing^{10–16}. These genes include the transcription factor *optix*^{14,17}, the signalling ligand *wntA*¹⁵ and the cell cycle regulator *cortex*¹⁶. Hence, these studies have revealed that a limited set of ‘toolkit’¹⁸ genes has been repeatedly used for both highly divergent and convergent phenotypes in *Heliconius*, as well as other butterfly and moth species^{16,19,20}. However, the key to wing pattern variation in *Heliconius* is not within the genes themselves, which are strongly conserved at the amino acid level, but at nearby non-coding regions that control expression during wing development^{14–16}.

Here, we sequenced the genomes of 15 distinctly coloured *H. erato* races and 8 closely related species to fully describe the regulatory architecture driving adaptive evolution of the major genes acting in *Heliconius* wing patterning (Fig. 1). Our genomic survey included samples obtained near seven transition zones of hybridizing *H. erato* races with divergent wing patterns (Fig. 2a). In these hybrid zones, the high rate of genetic admixture allows for detailed genotype by phenotype (G × P) association mapping to identify discrete genomic intervals associated with colour and pattern variation on *Heliconius* wings^{21,22}. We then further investigated these intervals with a novel phylogenetic method for identifying conserved non-coding regions in closely related non-hybridizing races and species. This combined strategy of association mapping and phylogenetic inference resulted in a distinct set of narrow genomic intervals that corresponded to loci described in early crossing experiments⁹

¹Department of Biology, Center for Applied Tropical Ecology and Conservation, University of Puerto Rico, Rio Piedras, Puerto Rico. ²Smithsonian Tropical Research Institute, Apartado 0843-03092, Panamá, Panama. ³Department of Zoology, University of Cambridge, Cambridge CB2 3EJ, UK.

⁴Hawkesbury Institute for the Environment, Western Sydney University, Richmond, New South Wales 2753, Australia. ⁵Biology Program, Faculty of Natural Sciences and Mathematics, Universidad del Rosario, Carrera. 24 No. 63C-69, Bogotá, DC 111221, Colombia. ⁶Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, Massachusetts, USA. ⁷Department of Ecology and Evolutionary Biology, Cornell University, 215 Tower Road, Ithaca, New York 14853-7202, USA. ⁸Department of Biology, Pennsylvania State University, University Park, Pennsylvania 16802, USA. ⁹PPG Biologia Animal, Departamento de Zoologia, Universidade Federal do Rio Grande do Sul, Av. Bento Gonçalves, 9500, Bloco IV, Prédio 43435, Porto Alegre, RS 91501-970, Brazil. ¹⁰Department of Biological Sciences, Mississippi State University, 295 Lee Boulevard, Mississippi 39762, USA. [†]These authors contributed equally to this work. [‡]These authors jointly supervised this work. *e-mail: vanbelleghemsteven@hotmail.com




Figure 1 | Geographical distribution, phylogeny and colour pattern diversity of the *Heliconius erato* adaptive radiation. a, Geographical origin of samples; colours represent the distribution of the races; numbers are placed according to the sampling sites. **b**, Maximum likelihood tree based on autosomal sites located on chromosomes that do not show any marked F_{ST} peaks. All nodes shown had full local support based on the Shimodaira–Hasegawa test. Colour and numbers represent the geographical distribution and sampling site, respectively. On average five individuals were sequenced for each race and two for each outgroup species. All samples used in this study were included in the tree. There were three cases (triangles) where individuals did not cluster together by racial designation (see Supplementary Fig. 5 for the full genome tree). **c**, Pictures of dorsal (left) and ventral (right) sides of the wings of races and species used in this study. Bottom row with black circles represent species that belong to the *erato* clade, but not to the *H. erato* adaptive radiation.

(Supplementary Table 1). All the intervals fell within non-coding regions adjacent to colour pattern genes that affect forewing band shape (*wntA*; Fig. 3), red pigmentation (*optix*; Fig. 4) and a yellow hindwing bar (*cortex*; Fig. 5). Our results underscore a highly modular regulatory architecture that provides a flexible mechanism for rapid morphological change (Fig. 6).

Results and discussion

Reference sequence and variants. With more than 25 different wing pattern races, *H. erato* provides exceptional opportunities to explore the links between genotype, phenotype, form and function. We first constructed a high-quality reference genome by a combination of hybrid assembly coupled with high-resolution linkage analysis. Our assembly and validation strategy generated one of the most contiguous and accurate Lepidopteran genomes assembled thus far (Supplementary Section 2), which is available on the LepBase genome browser. The final assembly consisted of 198 scaffolds with

N50 length of over 10 Mb and a total assembly length of 383 Mb. A total of 13,678 genes were identified using RNA-seq and a thorough annotation process (Supplementary Section 3). To examine variation across our reference genome, we generated high (15–30×) coverage whole-genome resequence data from 116 individuals of *H. erato* and closely related species. For the 101 *H. erato* individuals sampled, we genotyped the majority of the non-repetitive portion of the genome (average of 62% per individual; Supplementary Section 4.1). For the 15 individuals from the 8 outgroup species, the number of positions that were genotyped for the outgroup species was lower, but above 40% for the most divergent comparison (Supplementary Section 4.1).

Genome-wide divergence across the *H. erato* colour pattern radiation. Within *H. erato*, individuals clustered by geographic proximity rather than colour pattern phenotype, as has been previously reported²³ (Fig. 1b,c). For example, forewing red banded




Figure 2 | Genomic divergence across the *Heliconius erato* phenotypic transition zones. a, F_{ST} values were calculated between colour morphs from each of seven hybrid zones (indicated on the right) and averaged over 50 kb windows sliding in increments of 20 kb. Peaks represent regions of the genome with strongly divergent allele frequencies. Divergence at chromosomes 10, 15 and 18 corresponds with divergence near the colour pattern genes *wntA*, *cortex* and *optix* (red dashes), respectively. These loci drive black forewing, yellow hindwing bar and red pigmentation patterns, respectively. Importantly, between hybridizing races that were divergently coloured, the only regions of the genome in which we found fixed allelic differences were at the colour pattern loci (see Supplementary Section 4.3 for a discussion of other regions of the genome with increased divergence). **b**, Distribution of genotypes fixed between hybridizing races located in the peaks of high divergence. This analysis revealed that, depending on the variable phenotype in the hybrid zone, clusters of fixed SNPs are found in different genomic intervals near colour pattern genes.

H. erato races were found in all three (Caribbean/Pacific Coast, East Amazonian, and West Amazonian) major geographic lineages (Fig. 1). Even within these broad geographic regions, individuals used in this study grouped together by sampling location rather than wing morphology. Indeed, there was little genetic differentiation between *H. erato* individuals sampled across major phenotypic transition zones, except around the genomic regions already known to be involved in colour pattern variation (Fig. 2a). Genetic divergence as measured by F_{ST} (see Methods) was close to zero across most of the genome, supporting the hypothesis of unhindered gene flow except at the regions responsible for colour pattern differences ($F_{ST} < 0.1$ in $97.07 \pm 0.03\%$ of 50 kb windows; Supplementary Section 3.3)²². This contrasted with three sharp peaks of genomic differentiation across known colour pattern loci on chromosome 10 near the *wntA* gene, on chromosome 15 near *cortex*, and on

chromosome 18 near *optix* (shown in red in Fig. 2b). As previously reported for the region around *optix*²², these regions showed the expected signatures of selection, including reduced nucleotide diversity and elevated d_{XY} relative to genome-wide averages (Supplementary Section 4.3).

Associating genomic variation with colour pattern diversity. Genetic differences at the regions controlling phenotypic variation in *Heliconius* are maintained by strong natural selection^{24–26}. However, genotype by phenotype ($G \times P$) associations were often complex between any pairwise comparison reflecting different histories of interactions between hybridizing taxa. Thus, at any specific comparison, associations often spanned hundreds of thousands of base pairs around each colour pattern locus (Fig. 2b). Nonetheless, by combining analysis of variation across multiple hybrid zones




Figure 3 | Association mapping in hybrid zones and phylogenetic comparisons identify the modular genetic architecture of black forewing variation.

a, Variation in black forewing patterning in the *H. erato* races. Black shading in the forewings highlights variation in melanin production in different parts of the forewing. Colour shading corresponds to shading in **b** and **c**. **b**, F_{ST} (lines; 20 kb window, 5 kb step size) and association (points) analysis at the peaks of divergence in chromosomes 10 and 13. Coloured points represent associations estimated from fixed SNPs. **c**, Phylogenetic weighting of phenotypic hypothesis consistent with the Sd, St, Ly and Ro elements. These weightings were obtained by summing weightings for topologies that were consistent with the hypothesized groupings presented in the phylogenies. Tree topologies consistent with a geographic grouping are represented negative in grey. Within the genomic regions with high phylogenetic weighting support for a particular phenotypic hypothesis, we defined the boundaries of the colour pattern intervals as position 4,634,972–4,641,535 for Sd, 4,657,452–4,658,207 for St, 4,666,909–4,670,474 for Ly, and 4,700,932–4,708,441 for Ly, on chromosome 10 and position 14,341,251–14,412,364 for Ro on chromosome 13. It is possible to further subdivide the Sd interval into two narrow intervals based on the phylogenetic weighting support and patterns of shared genotypes (position 4,637,657–4,637,727 for Sd, and 4,639,853–4,641,535 for Sd). See Supplementary Sections 4.1.2 and 4.2.2 for the full phylogenetic trees of the identified intervals including all *H. erato* samples and closely related outgroup species.




Figure 4 | Modular architecture of red pattern variation. **a**, Variation in red colour patterning in the *H. erato* races in the ray (R), band (Y) and dennis (D) region of the wings. **b**, F_{ST} (lines; 20 kb window, 5 kb step size) and association (points) analysis at the peaks of divergence in the *optix* genomic region on chromosome 18 between races with red rays and dennis patch (ray-dennis) versus races with a red forewing band (postman) (red; top panel) and *H. e. amalfreda* (no rays) versus *H. e. erato* (rays) (brown; bottom panel). Coloured points represent associations estimated from fixed SNPs. **c**, Genotype weightings (10 SNP window, 5 SNP step size, 3 SNPs minimum genotyped in 50% of population) of the positions that were identified as fixed between ray-dennis versus postman. A weighting of 1 means races or species have the same genotypes as the postman races, whereas a weighting of 0 indicates completely different genotypes in the considered window of fixed SNPs. **d**, Phylogenetic weighting of phenotypic hypothesis consistent with the R, Y and D elements. These weightings were obtained by summing weightings for topologies that were consistent with the hypothesized groupings presented in the phylogenies. Due to haplotype sharing among rayed/dennis and postman races, tree topologies consistent with geography are never supported in this genomic interval. Support for topologies consistent with a geographic tree that accounts for this haplotype sharing are represented upside-down in gray. We outlined the following positions: 1,377,801–1,384,841 for R; 1,403,328–1,412,865 for Y; 1,420,912–1,422,355 for Y₂; 1,412,888–1,419,375 for D₁; and 1,422,585–1,428,307 for D₂ on chromosome 18. See Supplementary Section 3.3.2 for the full phylogenetic trees of the identified intervals including all *H. erato* samples and closely related outgroup species.




Figure 5 | Independent modules generate convergent yellow hindwing bar phenotypes. **a**, Variation in yellow hindwing bar in *H. e. favorinus* from Peru and *H. e. demophoon* from Panama. Note that the yellow hindwing bar morphology is not completely identical between these two races: the yellow hindwing bar of *H. e. demophoon* is narrow, long and pointing up, whereas *H. e. favorinus* exhibits a broader, shorter bar that points down. Shading corresponds to shading in **b**, where two independent association peaks are identified. **b**, F_{ST} (lines; 20 kb window, 5 kb step size) and association (points) analysis near the *cortex* gene on chromosome 15. Comparison between *H. e. favorinus* and *H. e. emma* (red) shows a block of divergence different from the comparison between *H. e. demophoon* and *H. e. hyudara* (green). The block of association between *H. e. demophoon* and *H. e. hyudara* overlaps with the *pam* gene, but no functional link with colour pattern variation has been identified for this gene¹⁶. Coloured points represent associations estimated from fixed SNPs. Based on fixed SNP associations, we defined the positions of these two intervals as 2,053,037–2,171,230 for Cr_1 (orange) and 2,211,881–2,315,926 for Cr_2 (yellow). See Supplementary Section 4.4.2 for the full phylogenetic trees of the identified intervals including all *H. erato* samples and closely related outgroup species.

with phylogenetic analysis, we pinpointed specific genomic intervals associated with specific aspects of phenotypic variation. This combination of $G \times P$ association and phylogenetic analysis revealed a highly modular architecture of the variation around major colour pattern loci.

Modular architecture of forewing black colour variation. Recent genetic mapping coupled with studies of gene expression, suggest that a single gene, *wntA*, is driving much of the forewing pattern variation across *Heliconius* species²⁷. Indeed, our $G \times P$ association highlighted a 100 kb non-coding region near *wntA* on chromosome 10 (Fig. 3). Clusters of fixed SNPs defined discrete genomic intervals associated with the phenotypic effects of the *Sd*, *St* and *Ly* loci that were first described more than 30 years ago⁹. Variation at *Sd*, *St* and *Ly* was predicted to control patterning across the middle to the most distal sections of the forewing, respectively (Fig. 3a). Consistent with this hypothesis, we identified: (1) a 25 kb region of fixed differences between *H. e. notabilis* and *H. e. lativitta* that differed across the lower (*Sd*) and the middle (*St*) region of the forewing (shown in purple in Fig. 3b); (2) a narrow peak of association

between *H. e. notabilis* and *H. e. etylus* that differed only in the lower forewing region (*Sd*) (blue in Fig. 3b); and (3) a broad region of association that spans roughly 60 kb and appears to be composed of several distinct peaks between *H. e. erato* and *H. e. hyudara* from French Guiana that differed in *St* and *Ly* (orange in Fig. 3b). Comparisons between races with identical forewings showed no $G \times P$ association across any of these regions (green in Fig. 3b).

To further refine the regions associated with forewing band pattern, we used a novel tree weighting approach called Twisst (topology weighting by iterative sampling of subtrees; see Methods)²⁸ to explore how phylogenetic relationships varied around *wntA*. We hypothesize that the genomic variation underlying wing pattern differences should cluster individuals by wing pattern rather than geographic proximity. Sliding window phylogenetic comparisons identified four narrow genomic intervals near *wntA* that were strongly associated with changes in the spatial distribution of black scales on the forewing (Fig. 3c). The first region was a 10 kb interval roughly 50 kb upstream of *wntA* (blue in Fig. 3c) that supported the monophyletic grouping of races that are partially black in the lower midsection of the forewing extending just distal of the discal cell

region. Similarly, a separate 8 kb interval roughly 35 kb upstream of *wntA* grouped geographically distant individuals with similar distribution of black scales across most of the distal mid-section of the forewing (*St* interval) (green in Fig. 3c). Finally, two additional regions, one 25 kb upstream of *wntA* and another centered on *wntA*, grouped all individuals that were partially black in the upper section of the forewing (*Ly* intervals) (orange in Fig. 3c). Although the region centered on *wntA* showed some support for tree topologies based on geographic proximity, we still considered it a possible colour pattern interval because the phenotypic grouping is more strongly supported than geographic grouping. Other areas across this region supporting the phenotypic tree also showed similar support for tree topologies based on geographic proximity and were not considered as candidate colour pattern intervals.

Our genomic analysis also confirmed a new locus (*Ro*) responsible for pattern variation in the most distal region of the forewing band²⁹. Comparisons of *H. e. notabilis* and *H. e. lativitta* showed an approximately 71 kb region associated with pattern differences in the upper forewing (purple in Fig. 3b). Similar to the *wntA* region, $G \times P$ associations were localized to non-genic regions near two genes, the *Heliconius* homologue of the *ventral veins lacking* gene (*vvl*) and the homologue of *radial spoke head protein 3* (*rsp3*). The transcription factor *vvl* is involved in the formation of specific wing veins, neuronal differentiation and steroid production in *Drosophila melanogaster*^{30–32}. The *rsp3* gene encodes a kinase-A-anchoring protein that scaffolds the cAMP-dependent protein kinase holoenzyme (PKA) and is involved in numerous regulatory events in the cell³³. The absence of geographically independent hybrid zones for this phenotype limited our ability to further resolve this region with phylogenetic weighting. Although spatial expression patterns of *wntA* in *Heliconius* have been shown to prefigure variation in this upper region of the forewing¹⁵, it is likely that one or both of these genes interact with *wntA* to shape this variation. Such epistatic interactions are commonly observed in colour pattern variation in *Heliconius*^{34–36}.

Modular architecture of red pattern variation. Regulation of red patterns across the fore- and hindwing of *H. erato*, known to be under control of the gene *optix*^{14,17}, was also highly modular. We identified discrete genomic intervals near *optix* that were associated with the presence of red hindwing rays, a red patch ('dennis') in the proximal part of the forewing and a red forewing band. We use the original nomenclature in *H. erato* for these different pattern elements: *R* for red hindwing 'rays', *D* for a red dennis forewing patch and *Y* for forewing 'band' colour (Fig. 4a)⁹.

Associations between individuals that differed across all three pattern elements, the so-called 'dennis-rayed' and 'postman' phenotypes, were strongly clustered in a 69 kb region downstream of *optix* (Fig. 4b)²⁶. Within this 69 kb region, $G \times P$ associations between hybridizing *H. e. amalfreda* and *H. e. erato*, which differ only by the absence/presence of hindwing rays, were clustered in a 7 kb interval (Fig. 4b). In this interval, *H. e. amalfreda* possessed the postman haplotype, which contrasts with the rest of the 69 kb region where *H. e. amalfreda* shared a haplotype with *H. e. erato*. Phylogenetic trees constructed from this region grouped *H. e. amalfreda* with postman phenotypes that lack rays (red shading in Fig. 4c). Unexpectedly, the tree across this interval clustered the outgroup species — *H. telesiphe*, *H. hortense*, *H. hecalesia*, *H. clysonymus*, and *H. sara* — on a derived node with all rayed *H. erato* races (Supplementary Section 5.3.2). *Heliconius hecalesia*, *H. hortense*, and *H. clysonymus* all have large red hindwing patches, whereas, *H. sara* and *H. telesiphe* possess much smaller red spots on the underside of their hindwing. This pattern contrasts with the phylogenetic placement of these species in the tree constructed with data from the rest of the genome (Fig. 1a), possibly reflecting historical introgression of modular elements among species closely




Figure 6 | Modular regulatory architecture characterizes colour pattern diversity within the *Heliconius erato* radiation.

The upper panel provides a summary of colour pattern variation found among *H. erato* butterflies that is related to spatial expression of the genes *wntA* (black forewing patterning; chromosome 10), *cortex* (yellow hindwing bar; chromosome 15), *optix* (red; chromosome 18) and a functionally uncharacterized genomic interval on chromosome 13 responsible for pattern variation in the most distal region of the forewing band (*Ro*; functional candidates *vvl* and *rsp3*). The boxes in the bottom panel represent chromosomal intervals that include regulatory modules. These regulatory modules are coloured for butterflies in which the pattern is expressed. The regulatory modules have been rearranged among *H. erato* races to generate distinct wing phenotypes. Note that for *Cr*₁ and *Cr*₂ and rays (*R*), band (*Y*) and dennis (*D*) patterns are expressed when, respectively, *cortex* and *optix* are expressed, whereas for *Sd*, *St* and *Ly* pattern expression corresponds with absence of *wntA* expression.

related to *H. erato*. Such patterns of introgression have also been observed in other closely related *Heliconius* species^{1,37}.

Genomic intervals strongly associated with forewing band colour (*Y*) and the red dennis patch (*D*) were similarly localized using the combination of $G \times P$ association and phylogenetic weighting. For forewing band colour, we identified two distinct and narrow intervals separated by approximately 20 kb (yellow in Fig. 4b,c). In these regions, there were 15 fixed SNPs that distinguished butterflies with a red forewing band from those that lacked red. Phylogenetic trees from this region strongly supported clustering of the red-banded phenotypes *H. telesiphe*, *H. hermathena*, *H. e. favorinus* and *H. e. hydara*, whereas *H. himera*, *H. hortense*, *H. clysonymus* and *H. hecalesia*, all of which lack red on the forewing, grouped with the yellow-banded *H. erato* races (Fig. 4c and

Supplementary Section 5.3.2). Finally, we identified several intervals associated with the red dennis patch. For this analysis, we focused primarily on genetic variation within *H. himera*. *Heliconius himera* has red on the hindwing similar to rays, but lacks the dennis patch. Therefore, comparing *H. himera* and *H. erato* races with a dennis/rays phenotype allowed us to separate the dennis from the ray elements. Across the 69 kb region, there was a 12 kb area where *H. himera* genotypes were similar to the postman haplotype (grey in Fig. 4b). Phylogenetic weighting analysis in this area strongly supported the grouping of *H. himera* individuals by colour pattern phenotype with postman races from both sides of the Amazon basin (grey in Fig. 4c).

Independent modules generate convergent yellow hindwing bar phenotypes. Recent association and expression data implicated the gene *cortex* as important in controlling a variety of pattern elements across the *Heliconius* wing, including presence or absence of a yellow hindwing bar in *H. erato*, known as the *Cr* locus^{9,16}. In *H. erato*, we identified two discrete regions containing clusters of fixed sites associated with a yellow hindwing bar in two geographically isolated, yet phenotypically similar, *H. erato* races (Fig. 5). The Peruvian races *H. e. favorinus* and *H. e. emma* differed across an interval consisting of 269 fixed SNPs over 100 kb roughly centered on *cortex* (red in Fig. 5). Eight of these SNPs fell within the coding region of *cortex*, but only one resulted in an amino acid substitution (an arginine to lysine at scaffold Herato1505 position 2,087,610). Curiously, a different region distinguished the Panamanian races *H. e. demophoon* and *H. e. hydata* (green in Fig. 5), which show a similar difference in the presence/absence of a yellow hindwing bar. In this hybrid zone, there was a cluster of fixed differences located roughly 100 kb away and centered on the *Heliconius* homologue of *parn*, a poly(A)-specific ribonuclease. These association differences are consistent with the independent evolution of the yellow hindwing bar on either side of the Andes^{34,38}.

In *H. erato*, there are other colour pattern elements controlled by variation at this locus, including the presence/absence of white hindwing fringes and a yellow forewing line³⁹, but our sampling of *H. erato* races did not allow us to distinguish these elements (Supplementary Section 5.4). The hybrid zone comparisons *H. e. notabilis*/*H. e. lativitta* and *H. e. notabilis*/*H. e. etylus* also showed increased F_{ST} estimates near the *cortex* gene, but no pattern of perfect association was observed for these comparisons. Crossing experiments have suggested possible epistatic interactions between *cortex* and *wntA*^{38,40}, which provides a possible explanation for this increased divergence without any phenotypic effect known to be directly controlled by the *cortex* locus. Furthermore, the phenotypic effects of alleles at this locus can be dramatic in other *Heliconius* species¹⁶, suggesting that this locus interacts broadly with the other *Heliconius* patterning loci^{10,41}.

Modular regulatory architecture and pattern diversity within *H. erato*. Less than 0.2% of the genome was associated with wing pattern diversity across the *H. erato* radiation. This variation was highly modular and fell in non-coding regions near colour patterning genes, including *optix*, *wntA* and *cortex*^{14–16}, as well as a less well-documented colour pattern locus (*Ro*) that controls spatial variation of melanin in the upper forewing. Based on the proximity of these mostly non-coding intervals to known patterning genes, it is likely they represent *cis*-regulatory regions modulating the spatial expression of key patterning genes in discrete areas of the developing wing. In *Heliconius*, this modularity of *cis*-regulatory architecture provides a mechanism for rapid evolution of novel morphologies.

Both shuffling of existing modules and *de novo* evolution of new modules is associated with phenotypic diversity in *H. erato*. Indeed, we can recreate the colour pattern diversity across the *H. erato* radiation

using a combination of non-genic regions near four colour pattern genes (Fig. 6). This conclusion is perhaps best exemplified in the distribution of genetic variation around *wntA*, where different colour pattern races have different combinations of four distinct genomic intervals. These different intervals are likely to regulate the expression of *wntA* in different areas of the forewing to adjust the position, size and shape of the forewing band to closely match patterns in other co-occurring warningly coloured butterfly species. Within this modular framework, recombination can reshuffle existing regulatory variation to generate new combinations of regulatory elements and new wing pattern phenotypes. Recombination of colour pattern modules and introgression into other populations is likely to be driven by high rates of gene flow between adjacent populations. For example, *H. e. amalfreda* appears to have evolved via recombination of regulatory variation between rayed (*H. e. erato*) and red-banded (*H. e. hydata*) haplotypes that instantaneously generated a novel wing pattern, a process that closely mirrors the one recently described in the co-mimetic forms of *H. melpomene*³⁷.

New regulatory modules associated with wing pattern variation can also evolve *de novo*, further increasing the flexibility of these regions to generate pattern diversity. This was evident in the independent evolution of the yellow hindwing bar in the *H. erato* clade (Fig. 5), and also in the comparison of regulatory variation around the red patterning locus between *H. erato* and its co-mimic *H. melpomene*. Red pattern variation in the two species is similarly generated by regulatory differences at the *optix* locus¹⁴, and the genomic position and order of its *cis*-regulatory elements is broadly similar²⁶. Furthermore, in both species distinct intervals were associated with different red pattern elements, and ‘enhancer shuffling’ through recombination has similarly generated novel red pattern phenotypes³⁷. This implies considerable conservation of function of *optix cis*-regulatory regions that were re-used to generate the convergent patterns that underlie mimicry. Nonetheless, the precise elements associated with placement of red in discrete areas of the fore- and hindwing are not homologous in the two species (Supplementary Section 5.3.3). Thus, convergent patterns are clearly independently derived in the two radiations by the parallel evolution of new enhancer variation.

Conclusion

Our results reconcile decades of genetic and genomic studies of *Heliconius* colour pattern variation^{9,42}. For the first time, we were able to place an entire radiation within a single genomic framework. This work has reinforced the role of a simple toolkit of a few colour pattern genes and demonstrated that pattern diversity is likely to be generated by the regulatory complexity around these genes. We have characterized a discrete number of 1–7 kb intervals that modulate phenotypic variation, and show that divergent and convergent morphologies, are the product of enhancer shuffling and *de novo* independent evolution of these modules. Overall, our work provides a genomic framework to further explore this regulatory complexity. The regions we identified may contain a number of distinct regulatory elements that may be further resolved with chromatin accessibility data⁴³ and studied in detail with targeted genome editing. Such an integrated genomic view promises to accelerate our understanding of the links between genotype and phenotype, and how they play out on a developing butterfly wing. This research has broader ramifications because the small number of genes shown to generate wing pattern variation across *Heliconius* have been implicated in pattern variation in other butterflies and moths^{16,19,44}. Thus, the *Heliconius* wing pattern loci appear to be ‘genomic hotspots’ that underlie the evolution of phenotypic diversity in Lepidoptera. The radiation of warning colours in *H. erato* provides an example of regulatory complexity generated by a small toolkit of genes. This may well be a common hallmark of rapid morphological diversification in adaptive radiations.

Methods

Scaffold assembly and validation. The *H. erato* (race *demophoon*) genome was assembled using Illumina paired-end reads with different insert sizes and partially gap filled with PacBio data (Supplementary Table 2). Illumina data was produced according to the ALLPATHS-LG assembly protocol⁴⁵ with the paired-end library originating from a single individual and the mate pair libraries from a second, sibling, individual. An initial assembly was performed with ALLPATHS-LG using default parameters and the reads were mapped back to the assembly to acquire accurate distributions of fragment size for each library. Next, contaminant small fragment sequences were purged from the paired-end and mate-pair libraries. Reads were error-corrected using the software Blue⁴⁶. A *kmer* database was built from the raw paired-end data and used to remove unsupported reads from mate-paired libraries. This step reduced polymorphism that may cause erroneous assembly. The PacBio data were error-corrected using the Illumina data and the LoRDEC software⁴⁷.

Five assemblies were obtained using different combinations of raw or error-corrected Illumina data. Each assembly was quality checked against approximately 4 Mb of BAC sequences using nucmer⁴⁸. All assemblies gave similar amounts of gapped sequence (about 10% of the base pairs), which reflects long simple repeats scattered across the genome. The assembly with the best statistics (that is, highest N50s and best alignment to BAC) was then post-processed to replace putative tandem repeats with Ns. Small repetitive scaffolds and putative redundant haplotype sequences were removed and based on a combination of 'all-versus-all' alignments and depth of coverage estimates prior to performing ALLPATHS-LG scaffolding. Gaps were then filled using the filled fragment pairs, the corrected PacBio data and the small scaffolds that had been previously removed using PBJelly⁴⁹. PBJelly was run three times iteratively to balance sensitivity and specificity and the final assembly, called *Hera_Stage1*, had a length of 402.8 Mb and scaffold N50 of 612 kb, respectively. The assembly process with associated statistics are provided in Supplementary Table 2 and Supplementary Fig. 1.

Linkage mapping. We generated a high-resolution linkage map by sequencing a backcross family generated from our focal genomic line (Supplementary Fig. 2). Our strategy was to identify markers by coupling high-coverage, whole-genome sequencing (30–40×) of each parent with low coverage (5–10×) sequencing of their offspring. The low sequencing coverage of the offspring makes it difficult to determine individual genotypes with high accuracy. We therefore developed an in-house pipeline utilizing the mpileup command in SAMtools⁵⁰ to produce genotype posteriors over a candidate set of 6.7 million SNPs. These genotype posteriors were used to construct a linkage map with Lep-Map3 (<https://sourceforge.net/projects/lep-map3/>), a new linkage mapping software developed from the Lep-Map1/2 software^{51,52}.

The linkage map was constructed with Lep-Map3 as follows (see Supplementary Figure 3 in SI section 2.3): First, to obtain the most accurate parent genotypes, we calculated the parental genotype posteriors using the combined information from parents and offspring using the ParentCall module (Lep-Map2). Next, we calculated pair-wise LOD scores between markers with zero recombination rate ($\theta = 0$) using the module SeparateIdenticals (Lep-Map3) with *lodLimit* = 26.5, *informativeMask* = 12 and *numParts* = 20. This step identified markers that segregated identically. The 20 most abundant identical maternal markers were used as the chromosome prints (each maternal marker in a chromosome segregates identically as there is no recombination in the female in *Heliconius* butterflies). In this step, we could identify 20 of the 21 chromosomes, because we found that chromosome 2 was completely homozygous in the mother. To identify chromosomes, especially chromosome 2, in the paternal linkage map, identical paternal markers were joined using module JoinLGs (Lep-Map3) with recombination rate $\theta = 0.01$ and LOD score limit *lodLimit* = 20. More precisely, the linkage groups could be linked together for chromosome 2 by inspecting the markers at nearby positions in the assembly. These paternal markers clustered to 21 linkage groups identifying chromosome 2 and the same 20 chromosomes that were found in the maternal map. Next, the module ShortPath (Lep-Map3) was run on the identical paternal markers. This module finds the longest shortest path in a marker graph (i.e. the longest path in a graph for which the shortest path is chosen between pairs of markers), where markers are nodes and each marker pair has been connected with an edge of length $4n - 3$, if there are n detected recombinations (different genotypes considering both phases in this case) between the markers. The best paths were manually checked to determine the final order of the markers. After the maternal and paternal markers were placed within a linkage framework (Supplementary Table 3), we added the remaining markers into this framework using JoinIdenticals (Lep-Map3), with LOD score limits of 25 and 20, for paternal and maternal markers, respectively. The 1.2 million markers that were heterozygous in both parents were discarded (*informativeMask* = 12). Finally, the identified linkage groups (chromosomes) were named to reflect the nomenclature of the *H. melpomene* genome. We were able to easily identify homologous chromosomes by mapping the flanking regions of each marker to the *H. melpomene* genome¹. Our final linkage map covered all 21 chromosomes, including the Z chromosome.

Assembly correction and chromosomal scaffolding. We used our high-resolution linkage map to error correct and improve our genome assembly. To do this, we first manually identified scaffolds that were inconsistent with our linkage map. About 10% of the scaffolds, representing 62 Mb, had such errors. Due to the high-density of markers on our linkage map, most errors were localized within a few kb. These errors generally fell at a gap sequence, meaning that the scaffolding step of the assembly process, rather than the creation of contigs, caused most misassemblies. The scaffolds in the assembly with errors were cut to produce an error-free assembly. The assembly was also separated into chromosomes at this point. There was about 16 Mb of gapped sequence in the assembly. The 34 scaffolds that failed to map to chromosomes totaled 3.7 Mb, 3.5 Mb of which were bacterial genome sequence and the rest was mainly very highly repetitive haplotypes that failed to create substantially long (>3 kb) contigs.

We produced the final assembly by integrating information from two independent *de novo* assemblies to gap fill our oriented stage2 assembly. The first was an ALLPATHS-LG assembly generated from the same Illumina dataset paired-end and mate-paired dataset, and assembled as follows. Illumina paired-end and mate-pair data were subsampled to prescribed coverage depth according to ref.⁴⁵ and assembled using ALLPATHS-LG with "HAPLOIDIFY = TRUE" and "CLOSE_UNIPATH_GAPS = False". The resulting assembly was improved by performing 3 iterations of PBJelly⁴⁹, incorporating prior PBJelly assemblies into subsequent iterations. The second was an assembly of an additional sibling female individual using approximately 100× coverage of 2 x 250 Illumina data generated from PCR free libraries. The genome of this individual was assembled using DISCOVAR *de novo*^{53,54}. The scaffolds that spanned gaps in our assembly were extracted from the BWA-MEM⁵⁵ produced bam files using in-house software. This software used a variant of Smith-Waterman local alignment⁵⁶ to compute the best alignment to fix gaps. Both positive and negative gaps were considered. The alignment parameters used were +1 for nucleotide match, -4 for mismatch, -8 for gap open and -1 for gap extension. Gaps were filled iteratively, using the independent ALLPATHS assembly first. Here we required an alignment score of 100 across a 4 kb region on each side of a gap for the gap to be filled. Regions with multiple gaps were joined as if they contained a single large gap. Finally, we filled remaining gaps using the DISCOVAR assembly. In this case, we used alignment to 2 kb regions around each gap. Using this strategy, we reduced the number of gaps in our assembly to 5.2 Mb. Assembly completeness, as assessed against a benchmarked set of 2,675 single-copy orthologues using BUSCO⁵⁷ was 82% (2,179) in the *H. erato* genome and a further 11% were present, but marked as 'fragmented'. These BUSCO results were similar to those for other high quality lepidopteran genomes (Supplementary Table 8). We assembled 5 of 20 autosomes and the Z chromosome into single scaffolds. We failed to identify a W chromosome, probably because of its highly repetitive nature. See Supplementary Figure 4 for the completeness of the scaffolding in the final *H. erato* genome assembly.

Genome annotation. Annotation of the genome was performed using Just_Annotate_My_Genome (JAMg; <https://github.com/genomecurator/JAMg>). To facilitate annotation, we used RNASeq data generated from different life stages and tissue types (Supplementary Table 9). These data include recent Illumina 2x250 data, 454 data, and archival Illumina 2x50 data. All data were preprocessed using 'justpreprocessmyreads' (<http://justpreprocessmyreads.sourceforge.net>) and were error corrected using Blue⁴⁶ with a 'reference' *kmer* dataset derived from the most recently collected 2x250 Illumina RNA-seq data and a coverage cut-off of 2. The Illumina RNA-Seq data was assembled using Trinity RNA-Seq version 2.1.1⁵⁸ with both the 'de-novo' and 'genome-guided' options. The 454 data alongside all mRNA data acquired from GenBank and public Illumina data acquired from NCBI SRA were assembled and clustered using MIRA 4.9.5⁵⁹. The Trinity *de-novo*, Trinity *genome-guided* and the MIRA assemblies were aligned and assembled against the genome using a new version of PASA⁶⁰, thus, creating a non-redundant, intron-aware transcript set referred here as PASA cDNA contigs. The new Illumina RNA-Seq were aligned against the reference *H. erato* genome using GSNAP v.2015-09-29⁶¹ providing high-quality information on intron coordinates. Repetitive content was identified (simple, complex/ transposable, *de novo*, tRNA and rRNA elements) using trf⁶², RepeatModeler⁶³, RepeatScout⁶⁴, RepeatMasker⁶⁵, RepBase data⁶⁵, tRNAScan⁶⁶ and Aragorn⁶⁷. This masked dataset was provided at the last stage of the pipeline only.

We used two *de novo* gene modellers, GeneMark-ET⁶⁸ and Augustus 3.2.1⁶⁹ for gene prediction. Both used the intron co-ordinates as external evidence. In addition, Augustus used further external evidence as hints including the RNA-seq coverage derived from the Illumina reads, protein domains acquired from searching the genome against Swissprot using the HHBlits program⁷⁰, a high-quality subset of the PASA cDNA contigs as determined by JAMg, alignments of Uniref50 and the *Heliconius melpomene* predicted protein set⁷¹. The Augustus HMM models were trained and evaluated using 'training' and 'test' subsets of the high-quality PASA cDNA contigs. Following this, the external evidence was weighted using the JAMg optimization method and the same training and test cDNA contig datasets. At this point, we determined that the repeat masking data provided inferior prediction results and thus they were not used in the final prediction. Finally, Augustus was run with UTR prediction enabled to reduce false positive exons. Resulting UTRs were removed from the final prediction.

The Repeat masking information, GenMark-ET, Augustus, PASA cDNA contigs, the Uniref50 and *H. melpomene* protein alignments were provided to EvidenceModeler⁷² to derive a consensus gene dataset. This consensus dataset was then twice edited with PASA2 in order to add alternative splicing information and the UTRs as supported by cDNA evidence. This formed our Official Gene Set (OGS1). The OGS1 proteins were then functionally annotated using Just_Annotate_My_Proteins (JAMp; <https://github.com/genomecurator/JAMp>) searched against Hidden Markov Profiles of known proteins with manually curated metadata (Swissprot; clustered at 70% identity and aligned). For each significant hit (using the default settings of JAMp such as an e-value of 1e-10 and p-value of 1e-12), any Gene Ontology, ENZYME and KEGG ontology terms of the known Swissprot proteins were linked to the *H. erato* predicted proteins but only if the annotation evidence was experimentally derived and not inferred (i.e. terms with the evidence codes of 'IEA', 'ISS', 'IEP', 'NAS', 'ND', 'NR' were ignored). The RNA-Seq data was finally aligned against the OGS1 CDS data and processed with DEW (<https://github.com/alpapan/DEW>) to infer the expression profiles for each gene. The functional and expression annotations are available from http://annotation.insectacentral.org/heliconius_erato.

Sequence alignment and variant calling. We collected and sequenced 101 individual *H. erato* butterflies from Peru ($n = 15$), French Guiana ($n = 14$), Suriname ($n = 5$), Ecuador ($n = 29$), Colombia ($n = 12$), Bolivia ($n = 4$), Mexico ($n = 6$) and Panama ($n = 16$). We collected phenotypically pure (i.e. phenotypes resembling the geographical *H. erato* races) individuals of each colour pattern race from admixed populations where the ranges of two colour pattern races overlap. Additionally, we collected individuals from 8 different closely related species including *H. ricini*, *H. sara*, *H. charithonia*, *H. hecalesia*, *H. telesiphe*, *H. hortense*, *H. clysonimus*, and *H. hermathena* (Fig. 1 and Supplementary Tables 10 and 11).

Whole-genome 100 bp paired-end Illumina resequencing data of these individuals was aligned to the *H. erato* v1 reference genome using BWA v0.7.13⁷³ with default parameters. PCR duplicated reads were removed using Picard v1.138 (<http://picard.sourceforge.net>) and sorted using SAMtools⁷⁴. Genotypes were called using the Genome Analysis Tool Kit (GATK) Haplotypecaller⁷⁵ with default parameters. Individual genomic VCF records (gVCF) were jointly genotyped using GATK's genotypeGVCFs with default parameters, except for setting expected heterozygosity to 0.025 to match the populations high heterozygosity and grouping individuals according to race and sampling location. Genotype calls were only considered in downstream analysis if they met the following criteria: quality (QUAL) ≥ 30 , minimum depth ≥ 10 , maximum depth ≤ 100 (to avoid false SNPs due to mapping in repetitive regions), overall depth $\leq 100 \times$ number of samples, strand bias (FS) < 200 , quality by depth ≥ 5 , and for variant calls, genotype quality (GQ) ≥ 30 .

Divergence and association analysis. We estimated levels of relative (F_{ST})⁷⁶ and absolute genetic divergence (d_{XY})⁷⁷, and nucleotide diversity (π)⁷⁷ between populations in sliding windows using python scripts and egglib⁷⁸. In all our analyses, we only considered windows for which at least 10% of the positions were genotyped for at least 75% of the individuals within each population. For the whole genome analysis of the seven hybrid zones, on average 96.4% (SD = 1.1%) of windows met these criteria. Genotype by phenotype (G \times P) associations were tested for each variant position using a two-tailed Fisher's exact test. Positions were excluded if less than 75% of individuals were genotyped for each phenotype. The sliding window approach and the identification of distinct blocks of associated SNPs provides a robust approach for identifying genomic regions of interests in our study system⁷⁹.

Phylogenetic analysis. We used FastTree v2.1⁸⁰ to infer an approximate maximum-likelihood phylogeny from the entire genome using the default parameters. In this analysis, we only used concatenated SNP data from chromosome 4–9, 11–14, 16, 17 and 20, because these chromosomes did not show any genetic divergence peaks in our population analysis. FastTree computes support values on nodes using the Shimodaira–Hasegawa test. Phylogenetic relationships of individuals across defined colour pattern intervals were constructed using maximum likelihood (ML) trees with RAxML v8.0.26⁸¹. The best likelihood tree was chosen from 100 trees generated from a distinct starting tree using a GTR model with CAT approximation of rate heterogeneity and the support values of this tree was inferred with 100 bootstrap replicates.

Phylogenetic weighting. We applied a phylogenetic strategy for identifying shared or conserved genomic intervals akin to 'phylogenetic shadowing'⁸². We evaluated the support for alternative phylogenetic hypotheses in the regions of peaks of divergence around colour pattern loci using a novel method called Twisst (topology weighting by iterative sampling of subtrees; <https://github.com/simonhmartin/twisst>)²⁸. This method solves the problem of describing the relationships between groups that are not necessarily monophyletic. Given a tree and a set of pre-defined groups (in this case races) Twisst determines a weighting for each possible topology describing the relationship of the groups (for example, 6 groups yield 105 possible unrooted topologies and therefore 105 weightings). Topology weightings are

determined by sampling a single member of each group and then identifying the topology matched by the resulting subtree. This sampling is iterated over a large number of subtrees and weightings are calculated as the frequency of occurrence of each topology. This method therefore reduces tree complexity caused by imperfect clustering of samples within groups. The ability to consider all possible topologies at each window provides an advantage over more commonly used likelihood ratio tests that only compare two topologies, which is especially relevant for taxa that have potentially many distinct evolutionary histories across their genomes. Weightings were estimated from 500 sampling iterations and averaged over ten bootstrap trees produced by RAxML v8.0.26⁸¹ for each 2 kb window. Averaging weightings over bootstrap trees is expected to reduce false support for certain phylogenetic groupings from trees with low bootstrap support.

For phylogenetic weighting along the *wntA* (chromosome 10) and *Ro* (chromosome 13) interval, we compared weightings of topologies defined by samples from the following six groups: *H. e. demophaon*, *H. e. etylus*, *H. e. notabilis*, *H. e. lativitta/emma*, *H. e. erato/amalfreda* and *H. e. hydara* (FG). To partly control for the strong phylogeographic signal within *H. erato*, we focused these analyses on eastern Andean and Amazonian races, which also show the most variation in forewing band shape, size and position. For the *optix* (chromosome 18) interval, we compared weightings of topologies defined by samples from the following six groups: *H. e. amalfreda*, *H. e. favorinus/hydara* (FG), *H. e. etylus/lativitta/emma/erato*, *H. himera*, *H. telesiphe* and *H. clysonymus/hortense/hecalesia*. To obtain weightings for hypothesized phylogenetic groupings of specific colour pattern forms, we summed the counts of all topologies that were consistent with the hypothesized grouping.

Genotype weighting *optix*. We evaluated genotypic similarity of species/races to the reference "postman" haplotype using a sliding window analysis. The "postman" haplotype was defined based on the consensus of fixed SNPs between all 'postman' (*H. e. demophaon*, *H. e. hydara* (Panama), *H. hydara* (French Guiana), *H. e. notabilis* and *H. e. favorinus*) and all 'rayed' (*H. e. erato*, *H. e. etylus*, *H. e. emma* and *H. e. lativitta*) *H. erato* races. In total there were 264 fixed SNPs across a 69 kb window on chromosome 18 near *optix*. For each species/race evaluated, the proportion of SNPs that were identical to the postman haplotype was calculated over windows of ten fixed SNPs, with a minimum coverage of 3 SNPs called in all individuals. The window size and minimum coverage was chosen to best capture the turn-over of the genotypic similarity along the genomic interval.

Defining boundaries of colour pattern intervals. Our argument for identifying regulatory modules was hierarchical. The association peaks, or regions of the genome containing clusters of sites perfectly associated with wing pattern phenotype, marked the genomic intervals that probably contained the functional variation responsible for phenotypic differences. We further resolved these intervals combining data across independent transition zones. The rationale is that independent recombination events in the distinct locations break down the pattern of associations, except at those very narrow intervals responsible for pattern differences. Thus, in these areas individuals should group by colour pattern phenotype rather than geographic proximity, which is the pattern evident across the bulk of the genome. This is the basis of the Twisst analyses described above. Specific boundaries are defined by a combination of Twisst and G \times P association. For example, near *wntA* and *optix*, we defined the boundary positions of the regulatory modules by overlaying the phylogenetic weighting with genotype tables of the fixed allelic differences in the hybrid zone comparisons. More precisely, at the regions where phylogenetic weighting support for phenotypic grouping shifted and increased rapidly, we conservatively identified the boundaries of the intervals by looking for patterns of shared genotypes between samples with similar phenotypes. It should be noted that this approach assumes a single origin for functional alleles that are shared across similar phenotypes and will miss regions where patterning alleles evolved independently. The boundaries of the regulatory modules near *Ro* and *cortex* were defined only using the fixed SNP associations because the geographic distribution of the phenotypes does not allow phylogenetic weighting to distinguish between geography and phenotypic grouping for these loci.

Data accessibility. Sequencing data was submitted to the Sequence Read Archive (SRA) with BioProject accession PRJNA324415; genome assembly data: SAMN05578372 to SAMN05578377; RNAseq data: SRR616674 to SRR616691, SAMN05578182 to SAMN05578206; linkage map data: SAMN05572290 to SAMN05572390; and re-sequencing data: SAMN05224096 to SAMN05224211.

Received 22 June 2016; accepted 13 December 2016; published 30 January 2017

References

1. Dasmahapatra, K. K. *et al.* Butterfly genome reveals promiscuous exchange of mimicry adaptations among species. *Nature* **487**, 94–98 (2012).
2. Lamichhaney, S. *et al.* Evolution of Darwin's finches and their beaks revealed by genome sequencing. *Nature* **518**, 371–375 (2015).

3. Brawand, D. *et al.* The genomic substrate for adaptive radiation in African cichlid fish. *Nature* **513**, 375–381 (2014).
4. Lamas, G. in *Hesperioidea – Papilionoidea*. Gainesville, Florida: Association for Tropical Lepidoptera (ed. Lamas, G.), 261–274 (Scientific Publisher, 2004).
5. Nijhout, H. F. *The Development and Evolution of Butterfly Wing Patterns*. (Smithsonian Institution, 1991).
6. Chouteau, M., Arias, M. & Joron, M. Warning signals are under positive frequency-dependent selection in nature. *Proc. Natl Acad. Sci. USA* **113**, 2164–2169 (2016).
7. Naisbit, R. E., Jiggins, C. D. & Mallet, J. Disruptive sexual selection against hybrids contributes to speciation between *Heliconius cydno* and *Heliconius melpomene*. *Proc. Biol. Sci.* **268**, 1849–1854 (2001).
8. Turner, J. R. G. A tale of two butterflies. *Nat. Hist.* **84**, 28–37 (1975).
9. Sheppard, P. M., Turner, J. R. G., Brown, K. S., Benson, W. W. & Singer, M. C. Genetics and the evolution of Müllerian mimicry in *Heliconius* butterflies. *Phil. Trans. R. Soc. B Biol. Sci.* **308**, 433–610 (1985).
10. Joron, M. *et al.* A conserved supergene locus controls colour pattern diversity in *Heliconius* butterflies. *PLoS Biol.* **4**, e303 (2006).
11. Papa, R. *et al.* Multi-allelic major effect genes interact with minor effect QTLs to control adaptive color pattern variation in *Heliconius erato*. *PLoS ONE* **8**, e57033 (2013).
12. Kronforst, M. R., Kapan, D. D. & Gilbert, L. E. Parallel genetic architecture of parallel adaptive radiations in mimetic *Heliconius* butterflies. *Genetics* **174**, 535–539 (2006).
13. Kapan, D. D. *et al.* Localization of müllerian mimicry genes on a dense linkage map of *Heliconius erato*. *Genetics* **173**, 735–757 (2006).
14. Reed, R. D. *et al.* *optix* drives the repeated convergent evolution of butterfly wing pattern mimicry. *Science* **333**, 1137–1141 (2011).
15. Martin, A. *et al.* Diversification of complex butterfly wing patterns by repeated regulatory evolution of a Wnt ligand. *Proc. Natl Acad. Sci. USA* **109**, 12632–12637 (2012).
16. Nadeau, N. *et al.* The gene *cortex* controls mimicry and crypsis in butterflies and moths. *Nature* **534**, 106–110 (2016).
17. Martin, A. *et al.* Multiple recent co-options of *Optix* associated with novel traits in adaptive butterfly wing radiations. *EvoDevo* **5**, 7 (2014).
18. Carroll, S. B. Evo-Devo and an expanding evolutionary synthesis: a genetic theory of morphological evolution. *Cell* **134**, 25–36 (2008).
19. Gallant, J. R. *et al.* Ancient homology underlies adaptive mimetic diversity across butterflies. *Nat. Commun.* **5**, 1–10 (2014).
20. Van't Hof, A. E. The industrial melanism mutation in British peppered moths is a transposable element. *Nature* **534**, 102–105 (2016).
21. Rosser, N., Dasmahapatra, K. K. & Mallet, J. Stable *Heliconius* butterfly hybrid zones are correlated with a local rainfall peak at the edge of the Amazon basin. *Evolution* **68**, 3470–3484 (2014).
22. Supple, M., Papa, R., Hines, H. M., McMillan, W. O. & Counterman, B. A. Divergence with gene flow across a speciation continuum of *Heliconius* butterflies. *BMC Evol. Biol.* **15**, 204 (2015).
23. Hines, H. M. *et al.* Wing patterning gene redefines the mimetic history of *Heliconius* butterflies. *Proc. Natl Acad. Sci. USA* **108**, 19666–19671 (2011).
24. Mallet, J. & Barton, N. H. Strong natural selection in a warning-color hybrid zone. *Evolution* **43**, 421–431 (1989).
25. Kapan, D. D. Three-butterfly system provides a field test of müllerian mimicry. *Nature* **409**, 18–20 (2001).
26. Supple, M. A. *et al.* Genomic architecture of adaptive color pattern divergence and convergence in *Heliconius* butterflies. *Genome Res.* **23**, 1248–1257 (2013).
27. Martin, A. *et al.* Diversification of complex butterfly wing patterns by repeated regulatory evolution of a Wnt ligand. *Proc. Natl Acad. Sci. USA* **109**, 12632–12637 (2012).
28. Martin, S. H. & Van Belleghem, S. M. Exploring evolutionary relationships across the genome using topology weighting. Preprint at *bioRxiv* <http://dx.doi.org/10.1101/069112> (2016).
29. Nadeau, N. J. *et al.* Population genomics of parallel hybrid zones in the mimetic butterflies, *H. melpomene* and *H. erato*. *Genome Res.* **24**, 1316–1333 (2014).
30. Danielsen, E. T. *et al.* Transcriptional control of steroid biosynthesis genes in the *Drosophila* prothoracic gland by *Ventral veins lacking* and *Knirps*. *PLoS Genet.* **10**, e1004343 (2014).
31. De Celis, J. F., Llimargas, M. & Casanova, J. *Ventral veinless*, the gene encoding the Cfla transcription factor, links positional information and cell differentiation during embryonic and imaginal development in *Drosophila melanogaster*. *Development* **121**, 3405–3416 (1995).
32. Meier, S., Sprecher, S. G., Reichert, H. & Hirth, F. *Ventral veins lacking* is required for specification of the tritocerebrum in embryonic brain development of *Drosophila*. *Mech. Dev.* **123**, 76–83 (2006).
33. Jivan, A., Earnest, S., Juang, Y.-C. & Cobb, M. H. Radial spoke protein 3 is a mammalian protein kinase A-anchoring protein that binds ERK1/2. *J. Biol. Chem.* **284**, 29437–29445 (2009).
34. Jiggins, C. D. & Mcmillan, W. O. The genetic basis of an adaptive radiation: warning colour in two *Heliconius* species. *Proc. R. Soc. B* **264**, 1167–1175 (1997).
35. Baxter, S. W., Johnston, S. E. & Jiggins, C. D. Butterfly speciation and the distribution of gene effect sizes fixed during adaptation. *Heredity* **102**, 57–65 (2009).
36. Huber, B. *et al.* Conservatism and novelty in the genetic architecture of adaptation in *Heliconius* butterflies. *Heredity* **114**, 515–524 (2015).
37. Wallbank, R. W. R. *et al.* Evolutionary novelty in a butterfly wing pattern through enhancer shuffling. *PLoS Biol.* **14**, e1002353 (2016).
38. Maroja, L. S., Alschuler, R., Mcmillan, W. O. & Jiggins, C. D. Partial complementarity of the mimetic yellow bar phenotype in *Heliconius* butterflies. *PLoS ONE* **7**, e48627 (2012).
39. Sheppard, P. M., Turner, J. R. G., Brown, K. S., Benson, W. W. & Singer, M. C. Genetics and the evolution of Müllerian mimicry in *Heliconius* butterflies. *Phil. Trans. R. Soc. B Biol. Sci.* **308**, 433–610 (1985).
40. Mallet, J. The genetics of warning colour in Peruvian hybrid zones of *Heliconius erato* and *H. melpomene*. *Proc. R. Soc. B* **236**, 163–185 (1989).
41. Joron, M. *et al.* Chromosomal rearrangements maintain a polymorphic supergene controlling butterfly mimicry. *Nature* **477**, 203–206 (2011).
42. Kronforst, M. R. & Papa, R. The functional basis of wing patterning in *Heliconius* butterflies: The molecules behind mimicry. *Genetics* **200**, 1–19 (2015).
43. Lewis, J. J. *et al.* ChIP-Seq-annotated *Heliconius erato* genome highlights patterns of cis-regulatory evolution in Lepidoptera. *Cell Rep.* **16**, 2855–2863 (2016).
44. Martin, A. & Reed, R. D. Wnt signaling underlies evolution and development of the butterfly wing pattern symmetry systems. *Dev. Biol.* **395**, 367–378 (2014).
45. Gnerre, S. *et al.* High-quality draft assemblies of mammalian genomes from massively parallel sequence data. *Proc. Natl Acad. Sci. USA* **108**, 1513–1518 (2011).
46. Greenfield, P., Duesing, K., Papanicolaou, A. & Bauer, D. C. Sequence analysis Blue: correcting sequencing errors using consensus and context. *Bioinformatics* **30**, 2723–2732 (2014).
47. Salmela, L. & Rivals, E. Sequence analysis LoRDEC: accurate and efficient long read error correction. *Bioinformatics* **30**, 3506–3514 (2014).
48. Kurtz, S. *et al.* Versatile and open software for comparing large genomes. *Genome Biol.* **5**, R12 (2004).
49. English, A. C. *et al.* Mind the gap: upgrading genomes with Pacific Biosciences RS long-read sequencing technology. *PLoS ONE* **7**, e47768 (2012).
50. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).
51. Rastas, P., Paulin, L., Hanski, I. & Lehtonen, R. Lep-MAP: fast and accurate linkage map construction for large SNP datasets. *Bioinformatics* **29**, 3128–3134 (2013).
52. Rastas, P., Calboli, F. C. F., Guo, B., Shikano, T. & Merilä, J. Construction of ultradense linkage maps with Lep-MAP2: Stickleback F2 recombinant crosses as an example. *Genome Biol. Evol.* **8**, 78–93 (2015).
53. Weisenfeld, N. I. *et al.* Comprehensive variation discovery in single human genomes. *Nat. Genet.* **46**, 1350–1355 (2014).
54. Love, R. R., Weisenfeld, N. I., Jaffe, D. B., Besansky, N. J. & Neafsey, D. E. Evaluation of DISCOVAR *de novo* using a mosquito sample for cost-effective short-read genome assembly. *BMC Genomics* **17**, 187 (2016).
55. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. Preprint at <https://arxiv.org/abs/1303.3997> (2013).
56. Smith, T. F. & Waterman, M. S. Identification of common molecular subsequences. *J. Mol. Biol.* **147**, 195–197 (1981).
57. Simão, F. A., Waterhouse, R. M., Ioannidis, P. & Kriventseva, E. V. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* **31**, 3210–3212 (2015).
58. Haas, B. J. *et al.* *De novo* transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nat. Protoc.* **8**, 1494–1512 (2013).
59. Chevreux, B. *et al.* Using the miraEST assembler for reliable and automated mRNA transcript assembly and SNP detection in sequenced ESTs. *Genome Res.* **14**, 1147–1159 (2004).
60. Haas, B. J. *et al.* Improving the *Arabidopsis* genome annotation using maximal transcript alignment assemblies. *Nucleic Acids Res.* **31**, 5654–5666 (2003).
61. Wu, T. D. & Nacu, S. Fast and SNP-tolerant detection of complex variants and splicing in short reads. *Bioinformatics* **26**, 873–881 (2010).
62. Benson, G. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids Res.* **27**, 573–580 (1999).
63. Smit, A. F. A., Hubley, R. & Green, P. RepeatMasker (2014); <http://www.repeatmasker.org/>
64. Price, A. L., Jones, N. C. & Pevzner, P. A. *De novo* identification of repeat families in large genomes. *Bioinformatics* **21**, 1351–1358 (2005).
65. Jurka, J. *et al.* Repbase Update, a database of eukaryotic repetitive elements. *Cytogenet. Genome Res.* **110**, 462–467 (2005).

66. Lowe, T. M. & Eddy, S. R. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.* **25**, 955–964 (1997).
67. Laslett, D. & Canback, B. ARAGORN, a program to detect tRNA genes and tmRNA genes in nucleotide sequences. *Nucleic Acids Res.* **32**, 11–16 (2004).
68. Lomsadze, A., Burns, P. D. & Borodovsky, M. Integration of mapped RNA-Seq reads into automatic training of eukaryotic gene finding algorithm. *Nucleic Acids Res.* **42**, e119 (2014).
69. Stanke, M., Schöffmann, O., Morgenstern, B. & Waack, S. Gene prediction in eukaryotes with a generalized hidden Markov model that uses hints from external sources. *BMC Bioinformatics* **11**, 1–11 (2006).
70. Remmert, M., Biegert, A., Hauser, A. & Johannes, S. HHblits: Lightning-fast iterative protein sequence searching by HMM-HMM alignment. *Nat. Methods* **9**, 173–175 (2012).
71. Davey, J. W. *et al.* Major improvements to the *Heliconius melpomene* genome assembly used to confirm 10 chromosome fusion events in 6 million years of butterfly evolution. *G3* **6**, 695–708 (2016).
72. Haas, B. J. *et al.* Automated eukaryotic gene structure annotation using EvidenceModeler and the Program to Assemble Spliced Alignments. *Genome Biol.* **9**, R7 (2008).
73. Li, H. & Durbin, R. Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics* **26**, 589–595 (2010).
74. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).
75. Van der Auwera, G. a. *et al.* From FastQ data to high-confidence variant calls: the Genome Analysis Toolkit best practices pipeline. *Curr. Protoc. Bioinform.* **11.10**, 1–33 (2013).
76. Hudson, R. R., Slatkin, M. & Maddison, W. P. Estimation of levels of gene flow from DNA sequence data. *Genetics* **132**, 583–589 (1992).
77. Nei, M. & Jin, L. Variances of the average numbers of nucleotide substitutions within and between populations. *Mol. Biol. Evol.* **6**, 290–300 (1989).
78. De Mita, S. & Siol, M. EggLib: processing, analysis and simulation tools for population genetics and genomics. *BMC Genet.* **13**, 27 (2012).
79. Nadeau, N. J. *et al.* Genomic islands of divergence in hybridizing *Heliconius* butterflies identified by large-scale targeted sequencing. *Phil. Trans. R. Soc. Lond. B. Biol. Sci.* **367**, 343–353 (2012).
80. Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2 – Approximately maximum-likelihood trees for large alignments. *PLoS ONE* **5**, e9490 (2010).
81. Stamatakis, A. RAXML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).
82. Bofelli, D. *et al.* Phylogenetic shadowing of primate sequences to find functional regions of the human genome. *Science* **299**, 1391–1394 (2003).

Acknowledgements

We thank A. Tapia for maintaining the *H. erato* genome line and for generating our mapping family, and M. Vargas and C. Rosales for Illumina library preparation. We acknowledge the University of Puerto Rico, the Puerto Rico INBRE grant P20 GM103475 from the National Institute for General Medical Sciences (NIGMS), a component of the National Institutes of Health (NIH); CNRS Nouragues and CEBA awards (B.A.C.); National Science Foundation awards DEB-1257839 (B.A.C.), DEB-1257689 (W.O.M.), DEB-1027019 (W.O.M.); awards 1010094 and 1002410 from the Experimental Program to Stimulate Competitive Research (EPSCoR) program of the National Science Foundation (NSF) for computational resources; and the Smithsonian Institution. This research was supported in part by Lilly Endowment, Inc., through its support for the Indiana University Pervasive Technology Institute, and in part by the Indiana METACyt Initiative. The Indiana METACyt Initiative at IU is also supported in part by Lilly Endowment, Inc.

Author contributions

S.M.V.B., B.A.C., W.O.M. and R.P. designed the study and wrote the paper. P.R., A.P. and J.J.M. conducted genome assembly. P.R. conducted linkage map and genome quality assessment. A.P. conducted genome annotation. S.M.V.B. conducted population genomic, phylogenetic and comparative genomic analyses. M.R., M.A.S., H.H. and J.J.H. conducted comparative genomic analyses. S.H.M. contributed scripts for Twist analyses. B.A.C., W.O.M., R.P., H.H., C.D.J., J.M., M.L., C.S., C.F.A. and G.M. collected samples for sequencing.

Additional information

Supplementary information is available for this paper.

Reprints and permissions information is available at www.nature.com/reprints.

Correspondence and requests for materials should be addressed to S.M.V.B.

How to cite this article: Van Belleghem, S. M. *et al.* Complex modular architecture around a simple toolkit of wing pattern genes. *Nat. Ecol. Evol.* **1**, 0052 (2017).

Competing interests

The authors declare no competing financial interests.